

بسم الله الرحمن الرحيم





شبكة المعلومات الجامعية التوثيق الالكتروني والميكرو فيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بعض الوثائق الأصلية تالفة





بالرسالة صفحات
لم ترد بالأصل





Role of MRI in Screening of Non-Traumatic Pediatrics Hip Pain

Thesis

Submitted for fulfillment of the Master degree
in Radio-diagnosis

By

Ghadier Ibrahim Attia

M.B.B.Ch

Faculty of Medicine-Ain Shams University

Under Supervision of

Prof. Dr./ Abeer Abd El Maksoud

Professor of Radio diagnosis

Faculty of Medicine – Ain Shams University

Dr./ Tarek Wahby

Lecturer of Radiodiagnosis

Faculty of Medicine – Ain Shams University

**Faculty of Medicine
Ain Shams University**

2020

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قالوا

لسبحناك لا علم لنا
إلا ما علمتنا إنك أنت
العليم العظيم

صدق الله العظيم

سورة البقرة الآية: ٣٢



Acknowledgments

First and foremost, I feel always indebted to **Allah**, the **Most Beneficent** and **Merciful** who gave me the strength to accomplish this work,

*My deepest gratitude to my supervisor, **Dr. Abeer Abd Elmaksoud**, Professor of Radiodiagnosis, Faculty of Medicine – Ain Shams University, for her valuable guidance and expert supervision, in addition to her great deal of support and encouragement. I really have the honor to complete this work under her supervision. God blesses her and keeps her a sun lightening the way for the scientific students.*

*I would like to express my great and deep appreciation and thanks to **Dr. Tarek Wahby**, Lecturer of Radiodiagnosis, Faculty of Medicine – Ain Shams University, for his meticulous supervision, his patience and granting me much of his time in reviewing and correcting this work.*

*Special thanks to my Parents and all my **Family** members for their continuous encouragement, enduring me and standing by me.*

Last but not least, I would also like to thank my colleagues, all members of Demerdash radio-diagnosis department and everyone helped me in this study.

 **Ghadier Ibrahim Attia**

List of Contents

<i>Subject</i>	<i>Page No.</i>
List of Tables.....	i
List of Figures	iii
Introduction	1
Aim of the Work.....	3
Review of Literature	
Anatomy of pediatric hip joint	4
Pathology of Different Lesions Causing Hip Pain.....	19
MRI Findings of Different Lesions Causing Hip Pain.....	45
Patients and Methods.....	67
Results.....	70
Illustrative cases	87
Discussion	95
Summary and Conclusion	99
References	101
Arabic Summary	—

List of Tables

<i>Table No.</i>	<i>Title</i>	<i>Page No.</i>
Table (1):	Number and percentage distribution of patients according to their sex (N=39).	70
Table (2):	Descriptive of patients according to their age (years) (N=39).	71
Table (3):	Number and percentage distribution of patients according to their side (N=39).	72
Table (4):	Number and percentage distribution of patients according to their MRI regarding Avascular necrosis (N=39).	73
Table (5):	Number and percentage distribution of patients according to their MRI regarding isolated hip effusion (N=39).	75
Table (6):	Number and percentage distribution of patients according to their MRI regarding synovitis (N=39).	76
Table (7):	Number and percentage distribution of patients according to their MRI regarding septic arthritis (N=39).	77
Table (8):	Number and percentage distribution of patients according to their MRI regarding Perthes (N=39).	78
Table (9):	Number and percentage distribution of patients according to their MRI regarding SCFE (N=39).	79

Table (10):	Number and percentage distribution of patients according to their MRI regarding osteomyelitis (N=39).....	80
Table (11):	Number and percentage distribution of patients according to their MRI regarding Osteomalacia (N=39).	81
Table (12):	Number and percentage distribution of patients according to their MRI regarding aneurysmal bone cyst (N=39).	82
Table (13):	Number and percentage distribution of patients according to their MRI regarding proximal femoral deficiency (N=39).	83
Table (14):	Number and percentage distribution of patients according to their MRI final diagnosis (N=39).	84
Table (15):	Relation between final diagnosis and demographic data (N=39).....	85

List of Figures

Figure No.	Title	Page No.
Figure (1):	Hip and Pelvis Anatomy	4
Figure (2):	The acetabulum.....	6
Figure (3):	The gross anatomy of the joint capsule	8
Figure (4):	Anterior ligament.....	9
Figure (5):	Illustrating the ischiofemoral ligament.....	10
Figure (6):	Normal hip joint of 10 years old child.....	11
Figure (7):	Sagittal image of the hip 12-year-old boy.	12
Figure (8):	Sagittal image of the hip 16-year-old boy. Note physis is up to close	12
Figure (9):	Coronal T2WI of 10 years old child showing normal acetabular labrum.....	13
Figure (10):	a&b: (a) T1 W coronal image depicts normal superior and inferior acetabular labrum (b) T2 W sagittal image of left hip showing anterior and posterior labrum which appear hypointense on both and are named according to location.....	13
Figure (11):	Normal hip joint of 10 years old child. Axial T1-weighted.	14
Figure (12):	Coronal T1-W GRE image of the right femoral head and acetabulum of a5-year- old boy.	15

- Figure (13):** Epiphyseal marrow transformation depicted on coronal T1-W images of the proximal femoral epiphysis 17
- Figure (14):** (a): Sagittal T2W FS image of left hip demonstrates the normal fibrous capsule appearing hypointense on all sequences and is reinforced by multiple ligaments. (b) T2W FS axial image shows iliopsoas tendon as a low intensity structure which is a major flexor of hip joint and is inserted on the lesser trochanter 17
- Figure (15):** a&b: (a) T2WFS Sagittal image of the left hip depicting normal transverse acetabular ligament and ligamentum teres. (b) T2W FS axial view depicting the normal ligamentum teres..... 18
- Figure (16):** SCFE-Kline's line 32
- Figure (17):** The lateral epiphyseal shaft angle measurement as described by Southwick on the frog lateral radiograph. 34
- Figure (18):** Types of proximal femoral focal deficiency..... 38
- Figure (19):** There is bowing of the femur, a lucent line perpendicular to the shaft (Looser's zone), and thickening of the medial femoral cortex in this patient with osteomalacia..... 40
- Figure (20):** Sagittal T1-weighted spoiled gradient-recalled echo fat-saturated contrast-enhanced MR image reveals central necrosis of right proximal femoral ossific

nucleus and sparing of anterior and posterior femoral head	46
Figure (21): Coronal T1-weighted spin-echo MR image shows femoral head signal hypointensity	47
Figure (22): Sagittal T1-weighted fast spin-echo MR image also reveals diffuse right proximal femoral ossific nucleus signal hypointensity and flattening	48
Figure (23): Coronal STIR MR image shows curvilinear right proximal femoral epiphyseal subchondral and right hip joint effusion signal hyperintensity).....	50
Figure (24): T1-weighted spoiled gradient-recalled echo fat-saturated contrast-enhanced subtraction MR image shows non enhancement of proximal femoral ossific nucleus central fragment confirming presence of necrosis.....	51
Figure (25): Coronal T1-weighted spin-echo MR image shows flattening and fragmentation of left proximal femoral ossific nucleus.....	52
Figure (26): MR T2-weighted, gradient- echo sequences, transversal): extensive joint effusion in the right hip	54
Figure (27): Coronal T2 weighted MR image of the hip joints in a 12-year-old child with JIA showing bilateral bright signal intensity within the joint space (arrows) which may represent synovial thickening or joint effusion. Focal bright signal is seen at the right femoral head due to bone erosion	55

- Figure (28):** Coronal T2 FAT SAT shows Moderate joint effusion. 56
- Figure (29):** Axial T1 FAT SAT with C image showing Loss of right femoro-acetabular joint space with extensive low T1/high fluid-sensitive signal within the acetabulum, the right femoral head and neck with focal areas of intramedullary contrast enhancement within the acetabulum and the femoral head..... 59
- Figure (30):** Radially reformatted MR images (double echo steady state sequence) with the femoral neck axis as the centre of rotation (shown here is the 1 o'clock position) depict the severity of posterior slippage, widening of the physis and joint effusion..... 60
- Figure (31):** Acute SCFE, associated bone marrow oedema is picked up by the STIR (short tau inversion recovery) sequence. Bone marrow oedema may be an early sign of avascular necrosis 61
- Figure (32):** MRI – Coronal section showing cartilaginous portion in PFFD 62
- Figure (33):** (A) Coronal T1-weighted magnetic resonance imaging demonstrating hypointense signal through the superiorlateral portion of the femoral neck consistent with a nondisplaced femoral neck fracture (arrow). (B) Coronal T2-weighted fat-suppressed magnetic resonance imaging showing corresponded increased signal intensity (long arrow) and significant joint effusion (short arrow). The