Safaa Mahmoud

بسم الله الرحمن الرحيم

مركز الشبكات وتكنولوجيا المعلومات قسم التوثيق الإلكتروني





Safaa Mahmoud



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأقراص المدمجة قد أعدت دون أية تغيرات





Potential role of hepatic stem/progenitor cells immunoexpression as a predictor to interferon response in Egyptian HCV patients

Thesis

Submitted for Partial Fulfillment of M.D. Degree in **Pathology**

By

Heba Ahmed Mahmoud Mohammed

M.B.B.CH, Msc. Pathology Faculty of Medicine, Ain- Shams University

Supervised by

Prof.Dr. Thanaa El-Sayed Ahmed Helal

Professor of Pathology Faculty of Medicine- Ain Shams University

Prof.Dr. Ahmed Mohy El -din Zaki

Professor of Pathology Faculty of Medicine- Ain Shams University

Prof.Dr. Ali Ahmed Ali Waheeb

Professor of Internal Medicine Faculty of Medicine - Al-Azhar University

Prof.Dr. Naglaa Samier Ahmed

Professor of Pathology Faculty of Medicine - Ain Shams University

Assist. Prof. Dr. Nehal Ahmed Radwan

Assistant Professor of Pathology Faculty of Medicine - Ain Shams University

Faculty of Medicine Ain Shams University 2019



Acknowledgments

First and foremost, I thank ALLAH for endowing me with health, patience, and knowledge to complete this work, as a part of generous help throughout my life.

Words stand short to express my deep respect and thanks to **Prof.Dr. Thanaa El-Sayed Ahmed Helal**, Professor of Pathology, Faculty of Medicine- Ain Shams University, for her great assistance, valuable advices and her endless help to accomplish this work. I really have the honor to work under her supervision.

I would like to extend my deepest thanks to **Prof.Dr. Ahmed Mohy El -din Zaki**, Professor of Pathology, Faculty of Medicine- Ain
Shams University, for his great support, fruitful advices and generous help throughout this work.

I wish to introduce my great respect and thanks to **Prof.Dr.** Ali Waheeb, Professor of Internal Medicine, Faculty of Medicine - Al-Azhar University, for the great help and suuport he has devoted to accomplish this work.

I would like to express my great appreciation and respect to **Prof.Dr. Naglaa Samier Ahmed,** Professor of Pathology, Faculty of Medicine - Ain Shams University, for her great outstanding support, unlimited help and valuable comments.

I am deeply thankful to Assist Prof. Dr. Nehal Ahmed Radwan, Assistant Professor of Pathology, Faculty of Medicine - Ain Shams University, for being there through all the work, for her sincere advices, unlimited help and endless encouragement.

I am really greatful to my **dear professors and colleagues** who have prayed for me, expressed concern and extended their hands.

To my family, my beloved father, my dear husband, and my sweet doughters. I am realy greatly indebted to you for providing me with love, encouragement and prayers throughout this work.

Meha Ahmed Mahmoud Mohammed

List of Contents

Subject	Page No.
List of Abbreviations	i
List of Tables	vi
List of Figures	viii
Introduction	1
Aim of the Work	4
Review of Literature	
Epidemiology and prevalence of viral hepatitis C	5
Risk factors of infection and mode of transmission	
Molecular Virology and pathogenetic mechanism of HC	V10
Immune responses and pathogenesis of hepatic injury in	
infections	
Clinical course and prognosis of HCV	
Histopathology of Chronic Virus C Hepatitis	
Pretratment requesits in interferone based therapy of chro	
Treatment	
Interferon based theraby	
Predictors of interferon treatment response HCV	
Hepatic progenitor/Stem cells	
Cytokeratin 19	
Neural cell adhesion molecule (CD56)	
Materials and Methods	
لمرجعية غير معرّفة.	خطأ! الإشارة ا
Discussion	- ,
Summary	
لمرجعية غير معرّفة.	
Recommendations	
References	
Arbic Summary	
This is summary	

List of Abbreviations

Abbr. Full-term

A : Activity grade based on the scoring system of

METAVIR

AFP : Alfa feto protein

Alb : Albumin

ALT : Alanine aminotransferase

BMI : Body mass index

VBT : Virological breakthrough

CAMs : Cell adhesion molecules

CC : Cholangiocarcinoma

CD : Cluster of differentiation

CK19 : Cytokeratine 19

CTL : Cytotoxic T-lymphocyte

DAB : Di-AminoBenzidine

DR : Ductular reaction

E : Envelope

ECM : Extracellular matrix

EDHS : Egyptian Demographic Health Survey

EDTA : Ethylene diamino tetraacetic acid

EFNA1 : A gene on chromosome 1q21-q22 that encodes

ephrin-A1 protein

EGF : Epithelial growth factor

EIAs : Enzyme immunoassays

EpCAM : Epithelial cell adhesion molecule

ETR : End of treatment response

EVR : Early virological response

F : Fibrosis Stage based on the scoring system of

METAVIR

FDA : Food and Drug Administration

FGF7 : Fibroblast growth factor 7

G0 : Mitotically quiescent state

G1-6 : Genotype 1-6

GGT : Gamma-Glutamyltransferase.

Hx&E: Hematoxylin and eosin

HA : Hepatocellular adenoma

HAI : Histologic activity index

HBV : Hepatitis B virus

HCC: Hepatocellular carcinoma

cHCC-CC: Combined hepatocellular-cholangiocarcinoma

HCV : Hepatitis C virus

HGF : Hepatic growth factor

HIV : Human immunodeficiency virus

HLA-C: Human leukocyte antigen C

HPCs : Hepatic stem/progenitor cells

HPF : High power field

IF : Intermediate filaments proteins

IFN : Interferon

IL-10R2 : Interleukin-10 receptor 2

IP-10 : IFN- γ -inducible protein

IPCs: Isolated progenitor cells

ISG : Interferon -stimulated genes

Jak-STAT: Janus kinase—signal transducers and

activators of transcription

LCC : Large cell change

METAVIR: Meta-analysis of Histological Data in Viral

Hepatitis

mRNA : Messenger Ribonucleic acid

NASH : Nonalcoholic steatohepatitis

NCAM : Neural cell adhesion molecule

NHANES: National Health and Nutrition Examination

Surveys

NK : Natural killer

NS : Non-structural

NVR : No virological response

PAS : Periodic acid-Schiff (stain)

PAT : Parenteral antischistosomal therapy

PBMCs : Nuclear factor-κB (NF-κB) peripheral blood

mononuclear cells

PBS: Phosphate Buffered Saline

P-C: Portacentral

PCR : Polymerase chain reaction

PDGF : Platlet drived growth factor

Peg-IFN: Pegylated interferon

P-P : Portal-portal

RBV : Ribavirin

RGT : Response-guided therapy

RNA : Ribonucleic acid

RVR : Rapid virological response

S : Steatosis degree according to Brunt's scale

SCC : Small cell change

SVR : Sustained virological response

TGF-β : Tissue growth factor- β

TH : T-helper lymphocyte

Thy-1: Thymus cell antigen-1

TIMP : Tissue inhibitors of metalloproteinases

TLR : Toll-like receptor

TMA : Transcription-mediated amplification

TNF-\alpha: Tumor necrosis factor-alpha

TWEAK: TNF-like weak inhibitor of apoptosis

USA : United States OF America

USP18 : Ubiquitin-specific peptidase 18

VBT : Virological breakthrough

VR : Virological relapse

WHO: World health Organization

Wnt : The Name Wnt is a portmanteau of int and Wg

and stands for "Wingless-related integration

site"

List of Tables

Table No	. Title
Table (1):	Interpretation of HCV Assays53
Table (2):	Algorithm for evaluation of the grades of activity according to the scoring system of METAVIR
Table (3):	American Association for the Study of Liver Diseases (AASLD) guide lines for Selection of Patients for Treatment
Table (4):	Summary of clinicopathological data of studied chronic HCV cases (no=120). أ! الإشارة المرجعية غير معرّفة.
Table (5):	Relation between virological response and demographic, laboratory and histopathologic parameters. خطأ! الإشارة المرجعية غير معرّفة.
Table (6):	Comparison between different HCV groups according to their response to treatment and their studied demographic, laboratory and histopathologic parameters. المرجعية غير معرّفة.
Table (7):	Median values of hepatic progenitor cells in the various groups of virological response. خطأ! الإشارة المرجعية غير معرّفة.
Table (8):	Hepatic progenitor cells expression by CK19 and CD56 (NCAM) in demographic, laboratory and pathologic variables in all studied chronic HCV cases. خطأ! الإشارة المرجعية غير معرّفة.

- Table (9): Relationship between hepatic progenitor cells expression by CK19 and CD56 (NCAM)and other demographic, laboratory and pathologic variables in all studied chronic HCV cases. خطأ! الإشارة المرجعية غير معرّفة.
- Table (10): Comparison between different grades of activity (A), different stages fibrosis (F) by METAVIR scoring and different degrees of steatosis (S) by Brunt's staging system system as regard the median of DR (CK19), the median of IPCs (CK19),the median of DR (CD56 (NCAM)) and the median of IPCs (CD56 (NCAM)) (Wilcoxan Rank Sum test). خطأ! الإشارة المرجعية غير معرّفة.

List of Figures

Figure No.	. Title
Figure (1):	Epidemic of hepatitis C virus (HCV) in the world
Figure (2):	HCV antibody prevalence (%) by major subdivision, EDHS 2008
Figure (3):	Genomic organization of hepatitis C polyprotein
Figure (4):	Schematic representation of the life cycle of HCV
Figure (5):	Components of Antiviral Immune Response
Figure (6):	Interferon signal transduction 16
Figure (7):	NK cell function in chronic hepatitis C infection
Figure (8):	Schematic diagram showing the pathway of extracellular matrix (ECM) production and degradation
Figure (9):	Schematic showing the four grades of portal-periportal activity (piecemeal necrosis) and confluent lobular necrosis according to Ishak's scoring system P-C, portal-central
Figure (10):	Schematic of the METAVIR algorithm 45
Figure (11):	Schematic showing METAVIR fibrosis stages F0, F1, F2, F3, and F446
Figure (12):	Structure of ribavirin

Figure (13):	Effects of RBV on innate and adaptive immune responses
Figure (14):	Graphic display of virological responses73
Figure (15):	Schematic overview of liver lobule82
Figure (16):	Liver stem cells in the canals of Hering, structures that connect bile ductules with parenchymal hepatocytes (bile duct and Hering canals are stained for cytokeratin 7)
Figure (17):	Mechanisms of HPCs differentiation during adult regeneration
Figure (18):	Pre-treatment HCV-RNA in the studied HCV cases (n=120). خطأ! الإشارة المرجعية غير معرّفة.
Figure (19):	Groups of HCV cases according to their response to treatment.خطأ! الإشارة المرجعية غير معرّفة.
Figure (20):	Grade of activity by METAVIR scoring system in the studied HCV cases (n=120). نارة المرجعية غير معرّفة.
Figure (21):	Stage of fibrosis by METAVIR scoring systemin the studied HCV cases (n=120)خطأ! الإشارة المرجعية غير معرّفة.
Figure (22):	Degree of steatosisin the studied HCV cases (n=120) خطأ! الإشارة المرجعية غير معرّفة.
Figure (23):	Median ductular reaction (DR) expression by CK19 in the various groups of virological response.خطأ! الإشارة المرجعية غير معرّفة.
Figure (24):	Comparison between responders and nonresponders regardingthe median ductular reaction (DR) expression by

- Figure (25): Median isolated progenitor cells (IPCs) expression by CK19 in the various groups of virological response.خطأ! الإشارة المرجعية غير معرّفة.
- Figure (26): Comparison between responders and nonresponders regardingthe median isolated progenitor cells (IPCs) expression by CK19 (Wilcoxn Rank Sum test). خطأ! الإشارة المرجعية غير معرّفة.
- Figure (27): Median ductular reaction (DR) expression by CD56 (NCAM) in the various groups of virological response. لإشارة المرجعية غير معرّفة.
- Figure (28): Comparison between responders and nonresponders regardingthe median ductular reaction (DR) expression by CD56 (NCAM) (Wilcoxn Rank Sum test). ثنارة المرجعية غير معرّفة.
- Figure (29): Median isolated progenitor cells (IPCs) expression by CD56 (NCAM) in the various groups of virological response. لإشارة المرجعية غير معرّفة.
- Figure (30): Comparison between responders and nonresponders regardingthe median isolated progenitor cells (IPCs) expression by CD56 (NCAM)(Wilcoxn Rank Sum test). خطأ! الإشارة المرجعية غير معرّفة
- Figure (31): Comparison between different grades of activity by METAVIR scoring system (A) as regard the median of DR (CK19) (I), the median of IPCs (CK19) (II), the median of DR (CD56 (NCAM)) (III) and the median of IPCs (CD56 (NCAM)) (IV). Capital letters represent the significance between groups (Wilcoxan Rank Sum test). خطأ! الإشارة المرجعية غير معرّفة.