



000000

تم رفع هذه الرسالة بواسطة / سنوي محمود عقل

بقسم التوثيق الإلكتروني بمركز الشبكات وتكنولوجيا المعلومات دون أدنى

مسئولية عن محتوى هذه الرسالة.

ملاحظات:





Comparative study between anterior and posterior approaches in management of cervical spondylotic myelopathy

Thesis

***Submitted for Partial Fulfillment of the Requirements for M.D degree in
Neurosurgery***

By

Ahmed Mohamed Elsayed Assar

M.B.B.Ch., M.Sc.

Supervised by

Prof. Dr. Esmail Hassan Sabry

Professor of Neurosurgery,
Faculty of Medicine - Ain Shams University

Prof. Dr. Hisham Anwer Abdelraheem

Professor of Neurosurgery
Faculty of Medicine - Ain Shams University

Prof. Dr. Emad Mamoun Hamza

Associate Professor of Neurosurgery
Faculty of Medicine - Ain Shams University

Dr. Sameh Mohammed Hefni

Lecturer of Neurosurgery
Faculty of Medicine - Ain Shams University

**Faculty of Medicine
Ain Shams University**



2021



بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَقَدْ نَزَّلْنَاهُ عَلَىٰ
عَلَمٍ مُّبِينٍ

صَدَقَ اللَّهُ الْعَظِيمُ



سورة طه - الآية (١١٤)

Acknowledgement

I'd like to thank Allah the mighty and the merciful for the blessings he granted me in my life.

I would like to express my deepest gratitude and appreciation to Prof. **Dr. Esmail Hassan Sabry**, Professor of Neurosurgery Ain Shams University, whose help, valuable directions, and objective criticism made the accomplishment of this work possible.

I am deeply indebted to **Prof. Dr.Hisham Anwar Abdelraheem**, Professor of Neurosurgery Cairo University, for his continuous encouragement, effort and his wide scientific vision.

I would also like to thank **Prof. Dr.Emad Mamoun Hamza**, Associate Professor of Neurosurgery Ain Shams University, who has provided me with a lot of material and has given me all the support needed to finish this work.

I am very grateful to **Dr. Sameh Mohammed Hefni**, Lecturer of Neurosurgery Ain Shams University, for his time, meticulous supervision and valuable advises without which this work would not have been accomplished.

I would finally like to thank my family, friends, and colleagues in Ain University for their continuous support and never ending will to help me accomplish this work.

Ahmed Mohamed Elsayed Assar

Dedication

I would like to dedicate this thesis to

To soul of my Father...

I miss you every single day

To all my family ...

Thank you for being my support system

I wouldn't be here without you

LISTOFCONTENTS

	<i>Page</i>
LIST OF TABLES	II
LIST OF FIGURES	IV
LIST OF ABBREVIATIONS.....	IX
INTRODUCTION.....	1
AIM OF THE WORK.....	3
Review of Literature	4
MATERIALS & METHODS.....	72
RESULTS.....	76
ILLUSTRATIVE CASES.....	112
DISCUSSION	124
SUMMARY AND CONCLUSION.....	132
REFERENCES	133
ARABIC SUMMARY	-

LIST OF TABLES

Table no.	Title	Page
1	Basic clinical data among 40 cervical spondylotic myelopathy patients.	78
2	Pre-operative neurological and radiological data among 40 cervical spondylotic myelopathy patients	79
3	Operative data among 40 cervical spondylotic myelopathy patients	80
4	Post-operative neurological and radiological data among 40 cervical spondylotic myelopathy patients	81
5	Post-operative outcome data among 40 cervical spondylotic myelopathy patients.....	82
6	Comparison between the 3 groups as regards basic clinical data using ANOVA and Chi square tests.....	85
7	Comparison between the 3 groups as regards pre-operative neurological and radiological data using ANOVA test	86
8	Comparison between the 3 groups as regards operative data using ANOVA test	89
9	Comparison between the 3 groups as regards post-operative neurological and radiological data using ANOVA test.....	90
10	Comparison between the 3 groups as regards post-operative outcome data using ANOVA and Chi square tests	92

Table no.	Title	Page
11	Comparison between anterior approach group of patients as regards serial neurological and radiological assessments	93
12	Comparison between posterior laminectomy with lateral fixation group of patients as regards serial neurological and radiological assessments	96
13	Pearson's correlation analysis for pre-operative Factors associated with recovery rate	104
14	Roc-curve of each technique to predict patient's recovery..	107
15	Roc-curve of each technique to predict patient's improvement	109

LIST OF FIGURES

Figure no.	Title	Page
1	Anatomy of first cervical vertebra, (a) superior surface, (b) inferior surface. (Wardak, 2012).....	6
2	Anatomy of second cervical vertebra, (a) superior surface, (b) inferior surface. (Combalia, 2013)	7
3	Occipital condyle anatomy (caudal view). (Combalia, 2013)	9
4	Internal ligaments of craniocervical junction. (Tubbs, 2011)	13
5	Lateral atlantooccipital ligament anatomy (b), (a) jugular foramen, (c) anterior longitunalligament, (d) rectus capitis lateralis muscle. (Tubbs, 2011).....	14
6	Lateral atlantooccipital ligament anatomy (b), (a) jugular foramen, (c) anterior longitudinal ligament, (d) rectus capitis lateralis muscle. (Tubbs, 2011).....	22
7	The mechanism of paradoxical movements of the atlas. The neutral position is the center figure. If the atlas is compressed anterior to the balance point, it flexes. If compressed behind the balance point, it extends. (both right and left figure) (Bogduk, 2000)	23
8	C1–C2 axial rotation. (a) the anterior arch of C1 glides around the odontoid process (arrows) (b) the lateral mass of C1 subluxates (arrow) forwards across the superior articular process of C2. (Ramieri, 2016)	24
9	Pillar view of upper cervical spine showing unique morphology of c2 and c2,3 joint orientation and transverse orientation of each level below c3. (Dowdell, 2018)	25

Figure no.	Title	Page
10	The instantaneous center of rotation (ICR) (Swartz, 2005)	26
11	Pathophysiological and biomechanical pathway of cervical spondylosis. (Ferrara, 2012)	28
12	Pathological change in degenerative cervical myelopathy patients. (Badhiwala, 2020).....	29
13	(a) normal cervical lordosis as dorsal vertebral body height less than ventral vertebral body height; (b) With degenerative process, loss of disc space height leads to loss of lordosis and elongation of moment arm; (d) applied to cervical spine and vertebral body compression; (c) Further elongation of moment arm leads to kyphosis postural. (Benzel, 2007)	30
14	Pathophysiology of degenerative cervical myelopathy. (Badhiwala, 2020).....	35
15	Modified Japanese Orthopaedic Association scoring system. (Badhiwala, 2020)	38
16	Nurick grades for myelopathy. (Badhiwala, 2020)	39
17	Anterior approach position the reverse Trendelenburg manner. (Cuzzocrea, 2019)	50
18	(A) Right sided anterior approach to the lower cervical spine after horizontal skin incision and longitudinal incision of the platysma. (B) With one retractor mobilizing the carotid sheath laterally and the other trachea and esophagus medially then the anterior surface of the cervical spine is exposed. (Klekamp, 2019).....	51

Figure no.	Title	Page
19	PLL removal by teasing in between the longitudinal fibers with a microcurette and a Kerrison can be used to remove. Beginning of removal of PLL is started the central portion of the PLL (Wiesel, 2013)	53
20	Limits of corpectomy, (A, B, C) the corpectomy should be done according to preoperative imaging from medial border of uncinates to other one and wide decompression at nerve root foramen. (Rhee, 2013).....	55
21	Types of cervical corpectomy, (a) partial corpectomy, (b) subtotal corpectomy, (c) multilevel subtotal corpectomy. (Pascal-Moussellard, 2019).....	57
22	Posterior laminectomy in both normal lordotic patient (a) and (b) in kyphotic patient, the cord does not migrate away from ventral compression. (Koh, 2014)	58
23	Prone position for posterior decompression. (Werner, 2015)	59
24	Open door laminoplasty, (a) the opening is done at lateral mass laminar junction with angled burr toward spinal cord not vertical to facet joint. (b) At opposite site, cut of lamina not completed leaving ventral bone intact. (c) greenstick fractures are created by placing dorso-lateral tension on the spinous process. (d) Ligamentum flavum removal and canal decompression are done. (Weinberg, 2020)	62
25	Steps of French door laminoplasty by spinous process-splitting and bilateral full thickness laminotomy at the lamina-facet junction then closure by interposition graft. (Girolami, 2017)	63
26	en-bloc laminectomy, (a) bilateral troughs are done by footplate from caudal to rostral with preservation of lateral mass, (b) removal the bone and thecal sac is decompressed (star). (Mueller, 2020).....	66

Figure no.	Title	Page
27	Skip laminectomy technique in four levels stenosis, (A, B, C) lateral and posterior aspects of the cervical spine illustrating C4 and C6 skip laminectomy for a consecutive four level decompression between C3–4 and C6–7 accomplished by removing alternate lamina (C4 and C6), the cephalad half of the C5 and C7 lamina and the ligamentum flavum at those four levels. (Shiraishi, 2002)	67
28	Roy Camille technique. (Vaccaro, 2017)	70
29	Different techniques of lateral mass screw insertion. (Mendoza, 2019)	71
30	Gender among 40 cervical spondylotic myelopathy patients	79
31	Post-operative outcome data among 40 cervical spondylotic myelopathy patients	84
32	Comparison between the 3 groups as regards duration of symptoms.....	86
33	Comparison between the 3 groups as regards pre-operative mJOA scale	87
34	Comparison between the 3 groups as regards preoperative MS scale.....	88
35	Comparison between the 3 groups as regards pre-operativecervical lordosis angle.....	88
36	Comparison between the 3 groups as regards pre-operative average C2-7 canal diameter.....	89
37	Comparison between the 3 groups as regards post-operativecervical lordosis angle.....	91

Figure no.	Title	Page
38	Comparison between the 3 groups as regards postoperative average C2-7 canal diameter	91
39	Comparison between pre- and post-operative mJOA scale assessment in anterior approach group	94
40	Comparison between pre- and post-operative MS scale assessment in anterior approach group	95
41	Comparison between pre- and post-operative Average C2-7 canal diameter assessment in anterior approach group	95
42	Comparison between pre- and post-operative mJOA scale assessment in posterior laminectomy with lateral fixation group	97
43	Comparison between pre- and post-operative MS scale assessment in posterior laminectomy with lateral fixation group	97
44	Comparison between pre- and post-operative mJOA scale assessment in posterior laminectomy alone group	99
45	Comparison between pre- and post-operative MS scale assessment in posterior laminectomy alone group	99
46	Comparison between pre- and post-operative average C2-7 canal diameter assessment in posterior laminectomy alone group.....	100
47	Comparison between the 3 groups of patients regarding serial mJOA scale assessments.....	101
48	Comparison between the 3 groups of patients regarding serial MS scale assessments	101

Figure no.	Title	Page
49	Comparison between the 3 groups of patients regarding serialcervical lordosis angle assessments	102
50	Comparison between the 3 groups of patients regarding serialaverage C2-7 canal diameter assessments	102
51	Correlation between duration of symptoms and recovery rate	105
52	Correlation between grade of muscle weakness and recovery rate	105
53	Correlation between mJOA scale and recovery rate	106
54	Correlation between MS scale and recovery rate.....	106
55	Correlation between number of operated cervical levels and recovery rate	107
56	ROC curve of anterior approach technique (patient's recovery)	108
57	ROC curve of posterior approach technique (patient's recovery)	109
58	ROC curve of anterior approach technique (patient's improvement)	110
59	ROC curve of posterior approach technique (patient's improvement)	111

LIST OF ABBREVIATIONS

[illegible]